

# Minimal residual disease monitoring by quantitative RT-PCR in core binding factor AML allows risk stratification and predicts relapse: results of the United Kingdom MRC AML-15 trial

John A. Liu Yin,<sup>1</sup> Michelle A. O'Brien,<sup>1</sup> Robert K. Hills,<sup>2</sup> Sarah B. Daly,<sup>1</sup> Keith Wheatley,<sup>3</sup> and Alan K. Burnett<sup>2</sup>

<sup>1</sup>Department of Haematology, Manchester Royal Infirmary, Manchester, United Kingdom; <sup>2</sup>Department of Haematology, Cardiff University School of Medicine, Cardiff, United Kingdom; and <sup>3</sup>Clinical Trials Unit, University of Birmingham, Birmingham, United Kingdom

The clinical value of serial minimal residual disease (MRD) monitoring in core binding factor (CBF) acute myeloid leukemia (AML) by quantitative RT-PCR was prospectively assessed in 278 patients [163 with t(8;21) and 115 with inv(16)] entered in the United Kingdom MRC AML 15 trial. CBF transcripts were normalized to  $10^5$  *ABL* copies. At remission, after course 1 induction chemotherapy, a  $> 3$  log reduction in *RUNX1-RUNX1T1* transcripts in BM in t(8;21) patients and a  $> 10$  *CBFB-*

*MYH11* copy number in peripheral blood (PB) in inv(16) patients were the most useful prognostic variables for relapse risk on multivariate analysis. MRD levels after consolidation (course 3) were also informative. During follow-up, cut-off MRD thresholds in BM and PB associated with a 100% relapse rate were identified: for t(8;21) patients BM  $> 500$  copies, PB  $> 100$  copies; for inv(16) patients, BM  $> 50$  copies and PB  $> 10$  copies. Rising MRD levels on serial monitoring accu-

rately predicted hematologic relapse. During follow-up, PB sampling was equally informative as BM for MRD detection. We conclude that MRD monitoring by quantitative RT-PCR at specific time points in CBF AML allows identification of patients at high risk of relapse and could now be incorporated in clinical trials to evaluate the role of risk directed/preemptive therapy. (*Blood*. 2012;120(14):2826-2835)

## Introduction

Despite improved survival rates, relapse remains the main cause of treatment failure in acute myeloid leukemia (AML). The core binding factor (CBF)-positive leukemias comprise AML with t(8;21) (q22;q22) and inv(16) (p13q22)/t(16,16) (p13;q22), which are characterized by the presence of *RUNX1-RUNX1T1* (*AML1-ETO*) and *CBFB-MYH11* fusion transcripts, respectively.<sup>1</sup> Although CBF AML belongs to the favorable cytogenetic subgroup, with a cure rate of  $> 65\%$  achievable with chemotherapy alone, in particular with the use of high-dose cytarabine-based consolidation regimens, disease relapse remains the most important single cause of treatment failure, occurring in up to 35% of patients.<sup>2-7</sup> At present, little is known about the kinetics of relapse or the development of resistance mechanisms; and furthermore, post-remission therapy in AML takes no account of the level of residual leukemia. However, the development of real-time quantitative RT-PCR techniques has enabled the sensitive detection and quantification of leukemia-associated genes as a measure of residual leukemia, thus allowing the tracking of reemergent leukemic clones. Fusion genes, such as *RUNX1-RUNX1T1* and *CBFB-MYH11* in CBF AML, *PML-RARA* in acute promyelocytic leukemia (APL), *DEK-CAN*, and mutations in the nucleophosmin (*NPM1*) gene and the Wilms' Tumor (*WT1*) gene, can now be used as molecular targets for monitoring residual leukemia.<sup>8-14</sup>

Monitoring of minimal residual disease (MRD) using quantitative RT-PCR has become an important diagnostic tool that permits the assessment of response to therapy and for detecting early

relapse during remission and to guide therapeutic decisions. Although MRD monitoring is now an integral part in the management of acute lymphoblastic leukemia<sup>15,16</sup> (generally done by multiparameter flow cytometry) and chronic myeloid leukemia,<sup>17</sup> its clinical utility in AML is currently restricted to the APL subtype, albeit where the relapse rate is now low, in which preemptive treatment at the time of molecular relapse has been shown to be of clinical benefit.<sup>11</sup> There is now cumulative evidence that monitoring of MRD can predict relapse in other subtypes of AML. In 2 recent studies from the German-Austrian AML Study Group, involving AML patients with inv(16)/t(16;16)<sup>18</sup> and *NPM1* gene mutations,<sup>19</sup> the authors reported clinically relevant MRD checkpoints that allowed the identification of patients who were at high risk of relapse. Early reports in AML with t(8;21) suggested that quantitative monitoring of MRD was useful in distinguishing patients at high risk of relapse from those in durable remission.<sup>20,21</sup> This observation was confirmed in subsequent studies in CBF AML using quantitative RT-PCR methodology.<sup>22-28</sup> However, in CBF AML, especially in the t(8;21) subtype, there has been, to date, a paucity of large prospective studies to assess the clinical utility of MRD monitoring, and in particular limited data on serial MRD assessment in BM and/or peripheral blood (PB) samples during and after chemotherapy.

Within the United Kingdom Medical Research Council AML-15 trial, we prospectively assessed the clinical value of serial MRD monitoring in CBF AML, and we herein report our findings.

Submitted June 7, 2012; accepted July 26, 2012. Prepublished online as *Blood* First Edition paper, August 8, 2012; DOI 10.1182/blood-2012-06-435669.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

The online version of this article contains a data supplement.

© 2012 by The American Society of Hematology

## Methods

### Patients and samples

The MRC AML-15 trial<sup>29</sup> compared 3 induction regimens (DA vs ADE vs FLAG-Ida), followed by randomization in consolidation (courses 3 and 4) to either MACE and MidAC or 2 doses of Ara-C (3 g/m<sup>2</sup> or 1.5 g/m<sup>2</sup>) or to stop or have a fifth course (Ara-C 1.5 g/m<sup>2</sup>). Patients were also randomized to receive gemtuzumab ozogamicin (GO; 3 mg/m<sup>2</sup>) at induction and/or consolidation in a 2 × 2 design. Allografting in the first complete remission (CR) was not recommended for CBF AML patients. At diagnosis, all trial patients had routine cytogenetic investigations and were also screened for the presence of *RUNX1-RUNX1T1* and *CBFB-MYH11* transcripts. The diagnosis of CBF AML was based on a positive cytogenetic and/or molecular result. Between July 2002 and January 2009, 361 CBF AML patients were recruited, of whom 278 [163 with t(8;21) and 115 inv(16)/t(16;16)], were included in the MRD study. BM and PB samples were requested at diagnosis, after each course of chemotherapy and serially every 3 months in the first year of follow up, every 4 months in the second year, 6 months in the third year, and at relapse. Informed consent was obtained from patients according to standard procedures, in accordance with the Declaration of Helsinki, in each center, and samples were collected as part of the treatment protocol, which was approved by the Wales Research Ethics Committee. Results of MRD monitoring were not disclosed to participating clinicians.

### Nucleic acid isolation, cDNA synthesis, quantitative RT-PCR

Methodologies for nucleic acid isolation and cDNA synthesis are described in supplemental Methods (available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article).

Real-time quantitative RT-PCR protocols for *RUNX1-RUNX1T1*, *CBFB-MYH11*, and *ABL* transcripts were performed on the ABI 7900HD platform, in line with recommendations of the EAC program.<sup>9,30</sup> For patients with inv(16), including t(16;16), only screening for the 3 major fusion subtypes A, D, and E comprising ~ 98% of all breakpoints was carried out. The absolute copy numbers of fusion gene transcripts was normalized to *ABL*, expressed per 10<sup>5</sup> copies of *ABL*. Median sensitivity of the quantitative RT-PCR assay for *RUNX1-RUNX1T1* and *CBFB-MYH11* transcripts was 10<sup>-5</sup> and was calculated as described.<sup>31</sup> PCR negativity and positivity were defined, according to EAC criteria.<sup>30</sup> Diagnostic and follow-up samples with < 10<sup>2</sup> and < 10<sup>3</sup> copies of *ABL*, respectively, were excluded from analysis.

### Clinical endpoints and statistical analysis

The outcome definitions (eg, RFS and OS) follow IWG guidelines.<sup>32</sup> CR was defined as < 5% blasts in a normocellular marrow, with peripheral neutrophil recovery to 1 × 10<sup>9</sup>/L, and platelet count to 100 × 10<sup>9</sup>/L, without evidence of extramedullary disease. Cumulative incidence of relapse (CIR) is measured from the date of achievement of remission (CR or CRi [without complete peripheral blood recovery]) until relapse with death as a competing risk. Survival from remission is measured from CR/CRi until death or date last seen.

### Statistical methods

Surviving patients were censored at January 1, 2009. Correlations were calculated using Spearman rank correlation coefficient, or Pearson correlation for log copy number as these showed approximate normality. Time-to-event outcomes are presented using Kaplan-Meier curves, with univariate analyses by the log-rank method. For multivariable analysis, Cox proportional hazards regression was used. To evaluate the effect of MRD positivity (as defined by transcript copy numbers) during sequential monitoring, Mantel-Byar analysis<sup>33</sup> was used to adjust for zero time-shift bias in that patients who relapsed after becoming MRD-positive will have been in remission while MRD-negative. In Mantel-Byar analysis, patients start in the MRD-negative group and transfer to the MRD-positive group when

**Table 1. Patient demographics and clinical characteristics**

	Total no.	t(8;21)	inv(16)
Patients	278	163	115
<b>Age, y</b>			
15-29	67	33	34
30-39	58	33	25
40-49	72	40	32
50-59	60	44	16
60+	21	13	8
Median (range)	42 (15-70)	45 (15-70)	38 (16-64)
<b>Sex</b>			
Female	118	69	49
Male	160	94	66
<b>Diagnosis</b>			
De novo	270	158	112
Secondary	8	5	3
<b>Performance Status (WHO)</b>			
0	198	120	78
1	67	39	28
2	6	2	4
3	5	2	3
4	5	0	2
<b>WBC, × 10<sup>9</sup>/L</b>			
< 10	99	76	23
10-19.9	49	39	10
20-49.9	67	36	31
50-99.9	35	10	25
100+	27	1	26
Missing	1	1	0
Median (range)	17.6 (1.2-298.0)	10.5 (1.2-153.0)	41.8 (1.3-298.0)

they become MRD-positive at the specified level. All outcomes are given at 5 years except where otherwise specified.

## Results

Patients' clinical characteristics are shown in Table 1; 163 patients with t(8;21) and 115 with inv(16) were included in the MRD study. Patients were excluded because of lack of or insufficient diagnostic and/or follow-up samples for molecular analysis; 8 patients were transplanted and were therefore censored from the study. All t(8;21) patients were positive for *RUNX1-RUNX1T1*, whereas 5 patients who were cytogenetically negative for inv(16)/t(16;16) were positive for the *CBFB-MYH11* transcript. The total number of samples analyzed in the study were as follows: 822 from BM and 855 from PB in t(8;21) patients and 681 from BM and 661 from PB in inv(16) patients. CBF transcript levels at diagnosis are shown in supplemental Results. Pretreatment *RUNX1-RUNX1T1* and *CBFB-MYH11* copy numbers in BM and PB did not correlate with age, WBC, performance status, secondary disease, or sex. Furthermore, pretreatment transcript levels in BM and PB for both t(8;21) and inv(16) patients, expressed as a log-transformed continuous variable, did not impact on CR, CIR, and OS in both unadjusted and adjusted analyses.

The median follow-up time for survival was 36 months (range, 2-79 months). The CR rates for all t(8;21) and inv(16) patients entered in the trial were 97% and 92%, with 5-year CIR of 18% and 23%, respectively. For the MRD study, data on 71 morphologically relapsed patients [38 t(8;21) and 35 inv(16)] were available for analysis. Comparison of CBF patients in the MRD study with those not in it (278 vs 83 not included) showed no difference in baseline characteristics, Mylotarg randomization, or RFS between the 2 groups.

**Table 2. Impact of BM and PB log reduction and log copy numbers at remission, on relapse risk: unadjusted and adjusted HRs per log decrease**


Measure	Unadjusted HR per log decrease	Adjusted HR per log decrease
<b>t(8;21)</b>		
Marrow log reduction	0.36 (0.18-0.74) <i>P</i> = .005	0.33 (0.15-0.73) <i>P</i> = .004
Marrow log copy level	2.27 (1.43-3.58) <i>P</i> = .0005	2.84 (1.61-5.00) <i>P</i> = .0002
Blood log reduction	0.70 (0.45-1.09) <i>P</i> = .12	0.65 (0.40-1.08) <i>P</i> = .09
Blood log copy level	1.46 (1.09-1.97) <i>P</i> = .01	1.57 (1.12-2.22) <i>P</i> = .007
<b>inv(16)</b>		
Marrow log reduction	0.33 (0.16-0.68) <i>P</i> = .003	0.30 (0.12-0.77) <i>P</i> = .02
Marrow log copy level	2.07 (1.20-3.58) <i>P</i> = .009	1.78 (0.98-3.21) <i>P</i> = .06
Blood log reduction	0.40 (0.21-0.76) <i>P</i> = .005	0.32 (0.15-0.71) <i>P</i> = .002
Blood log copy level	2.16 (1.39-3.34) <i>P</i> = .0006	2.37 (1.46-3.85) <i>P</i> = .0002

**Impact of log reduction and log copy numbers at remission after course 1 induction chemotherapy**

The impact of log reduction and log copy numbers (transcript levels) at remission on relapse risk was assessed by univariate and multivariate analyses in t(8;21) and inv(16) patients separately.


Multivariate analyses were adjusted for age, WBC, secondary AML, performance status, and sex.

**t(8;21).** For t(8;21) patients, the hazard ratios (HRs) according to relative log reduction in fusion transcripts and normalized level of *RUNX1-RUNX1T1* transcripts after induction are shown in Table 2. To determine a significant level for each of the 4 variables, further analysis was carried out on the BM and PB log reduction split in < 1, 1 or 2, 2 or 3, and > 3 groups and BM and PB transcript levels at cut-offs of 10, 50, 100, 500, and 1000 copies. The impact of BM log reduction on relapse risk is shown in Figure 1A; a > 3 log reduction is highly significant and is associated with a CIR of only 4% in the 47% of patients achieving it. With respect to BM copy numbers at remission, a level of < 100 copies identified 47% of patients with a CIR of 7% (representing 15% of all relapsing patients), whereas in PB, a cut-off level of 1000 copies was also prognostic, identifying 78% of patients with a CIR of 15%, compared with 50% in the remaining 22%. However, adjusted regression analysis using the 3 variables (BM log reduction and BM and PB copies) revealed that the BM log reduction was the most prognostic (*P* = .008). Adjusted analyses of survival from CR showed no significant effect of BM log reduction after



**Figure 1. Outcomes of log reduction in BM at remission in t(8;21) patients. P values are by log-rank test. (A) Cumulative incidence of relapse (CIR) by log reduction. (B) Survival from CR by log reduction.**

**Figure 2. Outcomes by *CBFB-MYH11* copy numbers in PB at remission in *inv(16)* patients.** *P* values are by log-rank test. (A) Cumulative incidence of relapse by copy numbers. (B) Survival from CR by copy numbers



induction (adjusted HR per log reduction = 1.09; 0.60-1.95; *P* = .8; Figure 1B).

***inv(16)*.** Similar analyses were carried out for patients with *inv(16)* at remission after course 1; the unadjusted and adjusted HR per log decrease are shown in Table 2. There was no critical BM log reduction level with respect to relapse risk, whereas other significant variables included BM copy numbers (> 100), PB log reduction (< 1 vs ≥ 1), and PB transcript levels. Of these, the most prognostic was the PB copy numbers, with a level of < 10 copies being associated with a CIR of 21% (Figure 2A), in the 51% of patients achieving it, representing 29% of all relapsing patients. Moreover, higher PB copy numbers had a significant adverse impact on survival after CR (adjusted HR per log reduction = 1.75; 1.04-2.94; *P* = .02; Figure 2B).

**Impact of log reduction and log copy numbers after courses 2 and 3**

Applying the same methodology as used for course 1, there was no additional prognostic value of BM and PB log reduction or log

copy numbers after course 2 in both groups of CBF AML than that already established after course 1.

***t(8;21)*.** After course 3, the 2 most prognostic factors for relapse risk were 4 log reduction in BM (< 4 log reduction, n = 20; CIR 49% vs > 4 log reduction, n = 18; CIR 13%; adjusted HR = 0.10; 0.01-0.83; *P* = .01) and BM copy number (copies > 500, n = 3; CIR 100% vs copies < 500, n = 55; CIR 28%; adjusted HR = 22.15; 3.75-130.76; *P* < .0001).

***inv(16)*.** The most prognostic factor after course 3 for relapse risk was a PB copy number of 10 (copies < 10, n = 36; CIR 36% vs copies > 10, n = 9; CIR 78%, adjusted HR = 3.07; 1.07-8.80; *P* = .03).

**Impact of GO on MRD at remission and consolidation**

The effect of GO, which we previously reported to be beneficial in the CBF subset,<sup>29</sup> on transcript levels after course 1 is shown in Table 3. Of 58 patients with *t(8;21)* log reduction data, 29 entered the GO induction randomization. There was some evidence that there were greater log reductions in transcripts in patients given GO

**Table 3. Effect of GO on transcript levels in CBF AML patients after course 1 of treatment**

	GO	No GO	P
t(8;21) < 1 log reduction	0	1	.04
t(8;21) 1-3 log reduction	5	10	
t(8;21) > 3 log reduction	9	4	
inv(16) < 10 copies	6	5	1.0
inv(16) 10-500 copies	3	5	
inv(16) > 500 copies	2	1	

(*P* = .04 for trend), but none for the 22 of 61 inv(16) patients entering GO randomization. After course 3, there was no difference in molecular response with respect to GO randomization in both subtypes (data not shown). When the effect of GO on relapse and survival was considered, the effect in this set of patients was consistent with the figures seen overall in the CBF leukemias.<sup>29</sup>

**Impact of chemotherapy regimens on MRD**

The numbers were small for a reliable comparison of chemotherapy regimens: FLAG-Ida appeared to provide a greater log reduction than DA or ADE after course 1 (data not shown), but this difference did not translate into significantly improved overall survival for FLAG-Ida patients.


**Impact of BM and PB copy numbers during follow-up**

The number of observations after course 4 was quite small (total of 82), and these data were therefore combined with those obtained


during the follow-up period, defined as > 4 weeks after completion of course 4. Serial monitoring in BM and PB was carried out until relapse or closure of the study. A key aim of serial monitoring during remission was to establish critical MRD thresholds in BM and PB to predict relapse. Relapse risk was assessed at different BM and PB cut-off transcript levels by Mantel-Byar analysis in both groups of CBF AML patients.

**t(8;21).** Mantel-Byar analysis of relapse after treatment identified that positivity at a rate of > 500 *RUNX1-RUNX1T1* copies was highly prognostic, identifying 15 of 78 patients with a high risk of relapse (Mantel-Byar estimates of relapse 100% vs 7%, *P* < .0001; Figure 3A). There was a significant difference in survival for these patients, with Mantel-Byar estimates of survival at 5 years of 94% vs 57% (HR = 40.18; 5.80-278.3; *P* = .001; Figure 3B). For PB, copy numbers of > 100 identified 15 patients of 86 with a Mantel-Byar estimate of relapse of 100%, compared with 7% in the MRD-negative group (*P* < .0001; Figure 3C) translating into survival from relapse of 59% vs 95% (*P* = .0007; Figure 3D). Of 127 remission patients, 11 had persistent MRD levels < 500 copies in the BM (ie, PCR positive), of whom 2 also had intermittent transcript levels < 50 copies in the PB; these patients have stayed in clinical remission with a median follow-up of 4.5 years (range, 2-8 years). Thus, in a small number of patients with t(8;21), low levels of MRD, as specified in this study, which did not increase on serial monitoring, were consistent with durable clinical remission.

**inv(16).** Patients with > 10 *CBFB-MYH11* copies in the BM had a Mantel-Byar estimate of relapse of 90%, which increased to 100% in the 17 of 53 patients who had copies of > 50, compared with 10% in patients with fewer than 50 copies (*P* < .0001; Figure



**Figure 3. Sequential MRD monitoring during follow-up in t(8;21) patients.** (A) Mantel-Byar estimate of relapse (%) in patients with > 500 *RUNX1-RUNX1T1* copies in BM. (B) Mantel-Byar estimate of survival (%) from CR in patients with > 500 *RUNX1-RUNX1T1* copies in BM. (C) Mantel-Byar estimate of relapse (%) in patients with > 100 *RUNX1-RUNX1T1* copies in PB. (D) Mantel-Byar estimate of survival (%) from CR in patients with > 100 *RUNX1-RUNX1T1* copies in PB.



**Figure 4. Sequential MRD monitoring during follow-up in inv(16) patients.** (A) Mantel-Byar estimate of relapse (%) in patients with > 50 *CBFB-MYH11* copies in BM. (B) Mantel-Byar estimate of survival (%) from CR in patients with > 50 *CBFB-MYH11* copies in BM. (C) Mantel-Byar estimate of relapse (%) in patients with > 10 *CBFB-MYH11* copies in PB. (D) Mantel-Byar estimate of survival (%) from CR in patients with > 10 *CBFB-MYH11* copies in PB.

4A). Survival from remission was 100% in those patients (n = 47) who did not reach the positive threshold (50 copies) compared with only 25% in patients (n = 17) who were positive (P < .0001; Figure 4B). Similarly, the 22 patients with > 10 copies in the PB had a Mantel-Byar estimate of relapse of 97%, compared with a relapse risk of 7% in those with < 10 copies (P < .0001; Figure 4C). This translated into survival from remission of 57% versus 91% (P = .0005; Figure 4D). Thus, in PB, any positive MRD level > 10 *CBFB-MYH11* copies was associated with clinical relapse.

**The evaluation of sequential quantitative RT-PCR monitoring to predict relapse**

The predictive value of sequential monitoring by quantitative RT-PCR in BM or PB could be demonstrated in 11 t(8;21) and 18 inv(16) patients who had rising MRD levels culminating in clinical relapse. The median times of detectable PCR positivity to hematologic relapse in BM and PB in t(8;21) patients were 4.9 months (range, 0.3-11 months) and 4.5 months (range, 2.3-7.3 months), respectively. Corresponding values for inv(16) patients were 3 months for both BM and PB (BM range, 1.3-9.6 months; PB range, 1.1-5.8 months). In the remaining 42 relapsed patients, inability to predict impending relapse was mostly the result of a lack of scheduled serial samples, as the intervals between BM or PB sampling for MRD analysis were > 3 months.


**Kinetics of relapse**

The kinetics of leukemia relapse in BM and PB in patients in whom serial monitoring data were available are shown in Figure 5. Relapse

kinetics varied between both patient and disease types. In t(8;21) patients, the median increments in *RUNX1-RUNX1T1* transcripts in BM and PB were 0.5 log/m (r = 0.2-1.5) and 0.6 log/m (r = 0.3-1.5), respectively, whereas in inv(16) patients, the corresponding median increment in *CBFB-MYH11* transcripts was 0.5 log/m in both BM and PB (BM: r = 0.1-1.7; PB: r = 0.0-2.5).

**Paired BM and PB MRD analysis**


To assess whether PB could be used for MRD monitoring, we analyzed paired BM and PB samples collected after induction (course 1), during consolidation (course 3), and during follow-up. For t(8;21) patients, there were 394 paired samples (81 after C1, 53 after C3, and 260 at follow-up) and for inv(16) patients, 369 paired samples (51 after C1, 61 after C3, and 257 at follow-up). With respect to *RUNX1-RUNX1T1* and *CBFB-MYH11* transcripts, the pairs were categorized into 4 groups as follows: PCR positivity in both BM and PB, PCR positive in BM and negative in PB, PCR positive in PB and negative in BM, and PCR negative in both BM and PB. The distribution in percentages among the groups at the various time points is shown in Figure 6. In both CBF AML subtypes, differences in transcript copies between BM and PB (ie, BM-positive and PB-negative) were larger during treatment and smaller during follow-up. The percentages of paired samples during follow-up belonging to the group with PCR positive in BM and PCR negative in the corresponding PB were 16% for t(8;21) and 12% for inv(16) patients, respectively. Conversely, there were very few samples that were PCR-positive in PB and PCR-negative in BM (2% in t(8;21) and 1% in inv(16) patients).



**Figure 5. Kinetics of relapse.** Graphs represent the rate of rise of normalized *RUNX1-RUNX1T1* and *CBFB-MYH11* transcript levels in serial samples before hematologic relapse. (A) t(8;21) BM. (B) t(8;21) PB. (C) inv(16) BM. (D) inv(16) PB.

We also compared transcript levels in paired BM and PB samples at diagnosis, after induction (C1), during consolidation (C3) and at first PCR positivity during follow-up. A significant

correlation between BM and PB MRD levels at these time points was found in both groups of patients, using both Spearman and Pearson analysis (Table 4).



**Figure 6. Quantitative RT-PCR analysis in paired BM and PB samples after induction, during consolidation, and at follow-up.** Corresponding pairs were categorized according to their quantitative RT-PCR values into 4 groups. The distribution in percentages among the groups at the 3 time points is illustrated by the columns. (A) t(8;21). (B) inv(16).

**Table 4. Correlation coefficients between paired BM and PB transcript copies at specific time points**

	Diagnosis	After 1 course	After 3 courses	After 4 courses and follow-up
t(8;21)	0.65	0.79	0.42	0.92
Spearman	$P < .0001$	$P < .0001$	$P = .005$	$P < .0001$
Log [t(8;21)]	0.64	0.78	0.50	0.74
Pearson	$P < .0001$	$P < .0001$	$P = .0005$	$P = .009$
inv(16)	0.40	0.45	0.33	0.95
Spearman	$P = .01$	$P = .007$	$P = .05$	$P < .0001$
Log [inv(16)]	0.17	0.49	0.44	0.84
Pearson	$P = .3$	$P = .0002$	$P = .01$	$P < .0001$

## Discussion

The aim of MRD monitoring in leukemias is to assess the effectiveness of treatment and to identify patients who are at high risk of relapse from those who are potentially cured. Specific gene targets can be accurately and reliably quantified using quantitative RT-PCR, thus facilitating the prospective clinical evaluation of MRD monitoring. However, in CBF AML, studies using quantitative RT-PCR techniques have been limited by relatively small numbers of patients and a lack of MRD data at specific time points during and after chemotherapy, especially in patients with t(8;21). Our MRD study in 163 t(8;21) and 115 inv(16) patients from a relatively young age group (median age, 42 years) represents, thus far, the largest cohort of CBF AML patients treated uniformly on the same chemotherapy regimens in the MRC AML-15 trial.

Several studies have shown that using nested RT-PCR in patients with t(8;21), *RUNX1-RUNX1T1* transcripts could be detected after chemotherapy, autologous and allogeneic bone marrow transplantation in many patients in long-term remission.<sup>34,35</sup> The phenomenon has been ascribed to quiescent populations of stem cells, monocytes and B cells harboring the fusion gene.<sup>36</sup> Similarly, in AML with inv(16), MRD studies using nested RT-PCR assays have reported persistence of residual disease in some long-term remitters, although most patients in prolonged remission remain PCR-negative.<sup>37-39</sup> Consequently, quantitative RT-PCR methods were developed to quantify *RUNX1-RUNX1T1* transcripts during remission in patients with t(8;21).<sup>20,21</sup>

Using a sensitive quantitative RT-PCR assay, we looked at relapse risk, with respect to both the reduction in leukemia load from diagnosis and the absolute residual disease levels, as measured by the *RUNX1-RUNX1T1* and *CBFB-MYH11* transcript copies, on achieving remission after course 1 induction chemotherapy. In t(8;21) patients, we showed that BM log reduction and BM (> 500 copies) and PB (> 1000 copies) were predictive of relapse risk, with a > 3 log reduction in bone marrow being the most prognostic variable on multivariate analysis. In inv(16) patients, no BM log reduction threshold for relapse risk could be established, but BM (> 100) and PB (> 10) copy numbers were predictive of which PB transcripts level was the most important prognostic variable. Assessment of MRD after course 3 (consolidation) was also shown to be informative. In t(8;21) patients, achievement of at least a 4 log reduction in fusion transcripts in the BM was associated with a CIR of only 10%, whereas conversely detection of MRD exceeding 500 *RUNX1-RUNX1T1* transcripts invariably predicted relapse. In the inv(16) group, PB MRD was most prognostic, with a *CBFB-MYH11* copy number > 10 being associated with a CIR of 78%.

We showed that, after consolidation chemotherapy and during remission follow-up, BM and PB transcript levels were highly predictive of relapse risk. In the t(8;21) group, the Mantel-Byar

estimates of relapse in patients with a BM *RUNX1-RUNX1T1* copy number > 500 (15 of 91 patients) or a PB copy number > 100 (15 of 99) were both 100%. Similarly, in inv(16), the Mantel-Byar estimate of relapse in patients with BM *CBFB-MYH11* transcript copies > 50 (16 of 77 patients) was 100%, whereas that in patients with PB copies > 10 (22 of 87) was 97%. Five-year survival in the MRD-positive patients in both t(8;21) and inv(16), as defined by transcript copy numbers, was significantly worse, compared with that in MRD-negative patients. However, it is worth noting that salvage rates in the relapsed t(8;21) patients were nearly 60%. In addition to demonstrating that critical MRD thresholds in both BM and PB could be established, above which relapse appears inevitable, we also showed that rising MRD levels on sequential monitoring during follow-up, which were not associated with BM or PB changes, accurately predicted hematologic relapse. It is also worth noting that in t(8;21) patients, low levels of MRD in both BM and PB were compatible with durable remission, consistent with the results previously described using qualitative RT-PCR. In contrast to inv(16) patients, molecular MRD negativity is not a prerequisite for long term remission in t(8;21) patients.

One of the aims of the AML-15 trial was to assess the effect of GO in AML patients, and we have previously reported that the addition of GO significantly improved the survival of CBF AML patients.<sup>29</sup> We therefore looked at MRD levels after courses 1 and 3 in these patients; there was some evidence that GO reduces molecular positivity after course 1 in t(8;21) but none for inv(16) patients. However, these results need to be interpreted with caution, as the number of evaluable patients is quite small. There was no difference in molecular responses after course 3 with respect to GO randomization. Marked molecular responses in CBF AML patients treated with GO and chemotherapy have been reported previously.<sup>40</sup> The results in our set of patients are in keeping with those seen overall in CBF AML.<sup>29</sup>

Several studies have evaluated the prognostic value of MRD monitoring by quantitative RT-PCR in t(8;21) patients undergoing chemotherapy.<sup>24-28</sup> In a study of 21 patients, Leroy et al reported that significant predictors of relapse were a > 3 log reduction in the BM and absolute transcript levels after induction and also high MRD levels after consolidation therapy.<sup>25</sup> Similarly Weisser et al studied 45 patients and showed that the quality of molecular response after both induction and consolidation chemotherapy was predictive of relapse risk, RFS, and OS.<sup>28</sup> In our study of 163 patients, we were able to confirm the prognostic value of BM log reduction and absolute MRD levels after both induction and consolidation chemotherapy. However, in our patients, although a > 3 log reduction after induction was prognostically important on multivariate analysis and predicted a relapse risk of only 4% compared with more than 30% for those who did not reach this threshold (HR per log reduction = 0.33; 0.15-0.73;  $P = .004$ ), it did not significantly impact on survival after CR, with a 5-year survival of 79%, which is no better than in the group with a 1-3 log reduction. This



observation, together with 5-year survival rates of nearly 60% in relapsed t(8;21) patients, can be explained by the fact that patients with CBF AML can be effectively salvaged after relapse.<sup>41</sup>

Similarly, the prognostic value of MRD monitoring by quantitative RT-PCR in inv(16) patients has been described in several studies.<sup>22,23</sup> Cut-off *CBFB-MYH11* copy numbers of 10-12 after completion of chemotherapy have been identified that could separate patients with high and low risk of relapse.<sup>27,42,43</sup> More recently, Corbacioglu et al, in a study of 53 *CBFB-MYH11*-positive AML patients, have reported clinically relevant MRD checkpoints during consolidation chemotherapy and early follow-up that allowed the identification of patients with high risk of relapse and also the negative impact of high MRD levels on RFS and OS.<sup>18</sup> They also showed that transcript levels below a threshold of 10 copies (normalized to  $\beta 2$ -microglobulin  $\times 10^6$ ) during follow-up were a prerequisite for long-term remission. In our study of 115 patients with inv(16), we were able to confirm that MRD levels based on *CBFB-MYH11* copy numbers after induction, during consolidation, and after chemotherapy were highly predictive of relapse risk. We also showed that a high MRD level in PB after induction adversely affected survival after CR.

One important aspect of this study was to assess the suitability of PB sampling compared with BM for MRD monitoring. Data on the applicability of PB for MRD assessment in CBF AML are sparse. In a small number of t(8;21) patients studied, a strong correlation between paired BM and PB MRD levels has been reported.<sup>25</sup> In inv(16) patients, a study by Boeckx et al on 54 paired BM/PB follow-up samples showed no clear correlation between BM and PB MRD levels, and these were generally higher in BM than in PB.<sup>44</sup> In contrast, Corbacioglu et al analyzed 198 paired BM and PB samples during chemotherapy and follow-up and found that 90% of the paired samples during follow-up showed either comparable or identical negative MRD levels.<sup>18</sup> In only 11% of samples, the PCR was positive in the BM, with a negative result in the corresponding PB. In our study, we analyzed 394 paired BM and PB samples in t(8;21) patients and 369 paired samples in inv(16) patients. In both groups of patients, differences in transcript copies between BM and PB (BM PCR-positive, PB PCR-negative group) were larger during treatment and smaller during follow-up, together with an increasing number of paired BM and PB samples with a negative PCR during follow-up. The percentage of paired samples with a positive PCR in the BM and negative PCR in the PB during follow-up in t(8;21) patients was 16%. In the corresponding group in inv(16) patients, the percentage was 12%. This result is very similar to that reported by Corbacioglu et al.<sup>18</sup> Interestingly, there were very few paired samples during follow-up that were PCR positive in PB and negative in BM (2% in t(8;21) and 1% in inv(16) patients). In addition, we showed that there was a significant correlation between BM and PB transcript levels at diagnosis, during treatment, and follow-up in both groups of CBF AML patients.

Information on the optimal schedules for MRD monitoring in CBF AML, including sampling source, is scanty, and published data are somewhat conflicting. The relapse kinetics in the 4 common AML subtypes with molecular targets (*NPM1c*, *PML-RARA*, *RUNX1-RUNX1T1*, and *CBFB-MYH11*) have all been shown to be strikingly different.<sup>11,18,19,26,45,46</sup> A study by Ommen et al applying a mathematical model to investigate optimal MRD monitoring by quantitative RT-PCR in CBF AML highlighted that *CBFB-MYH11*-positive AML relapses relatively more slowly than AML with *RUNX1-RUNX1T1*.<sup>46</sup> The authors proposed that, in inv(16) patients, PB monitoring every 6 months would be sufficient to predict relapse, whereas for t(8;21) patients, BM sampling every 4 months

would be required. In contrast, the recent study in *CBFB-MYH11*-positive AML by Corbacioglu et al showed that the kinetics of relapse are far more rapid than previously described, and our results accord with theirs.<sup>18</sup> We show that in inv(16) patients the kinetics of relapse in BM and PB were quite rapid with a median time from molecular positivity to clinical relapse of 3 months, whereas for t(8;21) patients, the median times to hematologic relapse were slightly longer (PB 4.5 months, BM 4.9 months) than in inv(16). Furthermore, in both t(8;21) and inv(16) patients, BM and PB were comparable for MRD detection during follow-up. We propose that optimal schedules for MRD monitoring in CBF AML should include assessment after induction, during consolidation, and at 3 monthly intervals in BM and/or PB, at least during the first 18 months of follow-up. Our data show that, during follow-up, MRD monitoring should ideally be carried out in paired BM and PB samples, to avoid missing a positive PCR result in BM with a negative PCR in the corresponding PB, in ~ 10%-15% of patients. However, in the majority of patients, PB sampling was equally informative for MRD monitoring and can therefore be used as a suitable alternative to BM.

In conclusion, MRD monitoring by quantitative RT-PCR in CBF AML allows risk stratification based on treatment responses after induction and consolidation chemotherapy, and sequential monitoring during follow-up can accurately predict relapse, thus opening the way to risk-directed therapy. Furthermore, we were able to establish MRD threshold levels in both BM and PB during follow-up, which were highly predictive of clinical relapse. In patients who have molecular relapse or rising MRD levels on serial monitoring, there may be a window of opportunity of ~ 3-4 months, which may allow for preemptive treatment, including investigational agents and potential allografting. Furthermore, patients who are *KIT* mutation positive may be candidates for a trial of tyrosine kinase inhibitors,<sup>47</sup> which, if successful, can also serve as a useful bridge to transplantation. Although monitoring of MRD in CBF AML may allow specific treatment to be tailored to individual patients, it is questionable whether a preemptive approach would be beneficial, given the rapidity of clinical relapse and the good responses achievable with salvage therapy. Thus, the role of early therapeutic intervention based on MRD detection to improve outcome in AML subtypes, other than APL, remains to be established in future clinical trials.

## Acknowledgments

The authors thank all consultant hematologists who participated in the AML 15 trial MRD study for providing diagnostic and follow-up samples and Dr David Grimwade for critical review of the manuscript.

This work was supported by the United Kingdom Leukemia and Lymphoma Research.

## Authorship

Contribution: J.A.L.Y., R.K.H., and A.K.B. initiated and designed the study, analyzed the data, and wrote the manuscript; M.A.O. and S.B.D. collected and analyzed the samples; K.W. designed the study; and all authors initially reviewed the final manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: John A. Liu Yin, Department of Haematology, Manchester Royal Infirmary, Oxford Road, Manchester, M13 9WL, United Kingdom; e-mail: john.yin@cmft.nhs.uk.

## References

- Swerdlow SH, Campo E, Harris NL, et al (eds). *WHO Classification of Haematopoietic and Lymphoid Tissues*. Lyon, France: IARC Press; 2008.
- Grimwade D, Walker H, Oliver F, et al. The importance of diagnostic cytogenetics on outcome in AML: analysis of 1,612 patients entered into the MRC AML 10 Trial. *Blood*. 1998;96(13):4075-4083.
- Byrd JC, Dodge RK, Carroll A, et al. Patients with t(8;21) (q22;q22) and acute myeloid leukemia have superior failure-free and overall survival when repetitive cycles of high-dose cytarabine are administered. *J Clin Oncol*. 1999;17(12):3767-3775.
- Byrd JC, Mrozek K, Dodge RK, et al. Pretreatment cytogenetic abnormalities are predictive of induction success, cumulative incidence of relapse and overall survival in adult patients with de novo acute myeloid leukemia: results from Cancer and Leukemia Group B (CALGB 8461). *Blood*. 2002;100(13):4325-4336.
- Byrd JC, Ruppert AS, Mrozek K, et al. Repetitive cycles of high-dose cytarabine benefit patients with acute myeloid leukemia and inv(16) (p13q22) or t(16;16): results from CALGB 8461. *J Clin Oncol*. 2004;22(6):1087-1094.
- Schlenk RF, Benner A, Krauter J, et al. Individual patient data-based meta-analysis of patients aged 16 to 60 years with core binding factor acute myeloid leukemia: a survey of the German Acute Myeloid Leukemia Intergroup. *J Clin Oncol*. 2004;22(18):3741-3750.
- Marcucci G, Mrozek K, Ruppert AS, et al. Prognostic factors and outcome of core binding factor acute myeloid leukemia patients with t(8;21) differ from those of patients with inv(16): a Cancer and Leukemia Group B Study. *J Clin Oncol*. 2005;23(24):5705-5717.
- Liu Yin JA, Grimwade D. Minimal residual disease evaluation in acute myeloid leukemia. *Lancet*. 2002;360:160-162.
- Gabert J, Beillard E, van der Velden VH, et al. Standardization and quality control studies of "real-time" quantitative reverse transcriptase polymerase chain reaction of fusion gene transcripts for residual disease detection in leukemia: a Europe against cancer program. *Leukemia*. 2003;17(12):2318-2357.
- Schnittger S, Weissner M, Schoch C, Hiddemann W, Halerlach T, Kern W. New score predicting for prognosis in PML-RARA, AML1-ETO, or CBFβ-MYH11 acute myeloid leukemia based on quantification of fusion transcripts. *Blood*. 2003;102(8):2746-2755.
- Grimwade D, Jovanovic JV, Hills RK, et al. Prospective minimal residual disease monitoring to predict relapse of acute promyelocytic leukemia and to direct pre-emptive arsenic trioxide therapy. *J Clin Oncol*. 2009;27(22):3650-3658.
- Tobal K, Frost L, Liu Yin JA. Quantification of DEK-CAN fusion transcript by real-time reverse transcription polymerase chain reaction in patients with t(6;9) acute myeloid leukemia. *Haematologica*. 2004;89(10):1267-1269.
- Cilloni D, Gottardi E, De Micheli D, et al. Quantitative assessment of WT1 expression by real-time quantitative PCR may be a useful tool for monitoring minimal residual disease in acute leukemia patients. *Leukemia*. 2002;16(10):2115-2121.
- Grimwade D, Vyas P, Freeman S. Assessment of minimal residual disease in acute myeloid leukemia. *Curr Opin Oncol*. 2010;22(6):656-663.
- Brüggermann M, Raff T, Flohr T, et al. Clinical significance of minimal residual disease quantification in adult patients with standard-risk acute lymphoblastic leukemia. *Blood*. 2006;107(3):1116-1123.
- Borowitz MJ, Devidas M, Hunger SP, et al. Clinical significance of minimal residual disease in childhood acute lymphoblastic leukemia and its relationship in other prognostic factors: a Children's Oncology Group Study. *Blood*. 2008;111(12):5477-5485.
- Müller MC, Cross NC, Erben P, et al. Harmonisation of molecular monitoring of CML therapy in Europe. *Leukemia*. 2009;23(11):1957-1963.
- Corbacioglu A, Scholl C, Schlenk RF, et al. Prognostic impact of minimal residual disease in CBFβ-MYH11-positive acute myeloid leukemia. *J Clin Oncol*. 2010;28(23):3724-3729.
- Krönke J, Schlenk RF, Jensen KO, et al. Monitoring of minimal residual disease in NPM1-mutated acute myeloid leukemia: a study from the German-Austrian Acute Myeloid Leukemia Study Group. *J Clin Oncol*. 2011;29(19):2709-2716.
- Tobal K, Liu Yin JA. Monitoring of minimal residual disease by quantitative reverse transcriptase polymerase chain reaction for AML-MTG8 transcripts in AML-M2 with t(8;21). *Blood*. 1996;88(10):3704-3709.
- Tobal K, Newton J, Macheta M, et al. Molecular quantification of minimal residual disease in acute myeloid leukemia with t(8;21) can identify patients in durable remission and predict clinical relapse. *Blood*. 2000;95(3):815-819.
- Buonamici S, Ottaviani E, Testoni N, et al. Real-time quantification of minimal residual disease in inv(16) positive acute myeloid leukemia may indicate risk for clinical relapse and may identify patients in a curable state. *Blood*. 2002;99(2):443-449.
- Guerrasio A, Pilatino C, De Micheli, et al. Assessment of minimal residual disease (MRD) in CBFβ-MYH11 positive acute myeloid leukemia by quantitative RT-PCR amplification of fusion transcripts. *Leukemia*. 2002;16(6):1176-1181.
- Krauter J, Gorlich K, Ottmann O, et al. Prognostic value of minimal residual disease quantification by real-time reverse transcriptase polymerase chain reaction in patients with core binding factor leukemias. *J Clin Oncol*. 2003;21(23):4413-4422.
- Leroy H, de Botton S, Grardel-Duflos N, et al. Prognostic value of real-time quantitative PCR (RQ-PCR) in AML with t(8;21). *Leukemia*. 2005;19(3):367-372.
- Stentoft J, Hokland P, Ostergaard M, Hasle H, Nyvold CG. Minimal residual core binding factor AMLs by real time quantitative PCR: initial response to chemotherapy predicts event free survival and close monitoring of peripheral blood unravels the kinetics of relapse. *Leuk Res*. 2006;30(4):389-395.
- Perea G, Lasa A, Avenirin A, et al. Prognostic value of minimal residual disease (MRD) in acute myeloid leukemia (AML) with favourable cytogenetics (t(8;21) and inv(16)). *Leukemia*. 2006;20(1):87-94.
- Weisser Haferlach C, Hiddemann W, Schnittger S. The quality of molecular response to chemotherapy is predictive for the outcome of AML1-ETO positive AML and independent of pretreatment risk factors. *Leukemia*. 2007;21(16):1177-1182.
- Burnett AK, Hills RK, Milligan D, et al. Identification of patients with acute myeloblastic leukaemia who benefit from the addition of Gemtuzumab Ozogamicin: results of the MRC AML 15 Trial. *J Clin Oncol*. 2011;29(4):369-377.
- Beillard E, Pallsigaard N, van der Velden VH, et al. Evaluation of candidate control genes for diagnosis and residual disease detection in leukemic patients using "real-time" quantitative reverse-transcriptase polymerase chain reaction (RQ-PCR): a Europe Against Cancer Program. *Leukemia*. 2003;17(12):2474-2486.
- Freeman SD, Jovanovic JV, Grimwade D. Development of minimal residual disease-directed therapy in acute myeloid leukemia. *Semin Oncol*. 2008;35(4):388-400.
- Cheson BD, Bennett JM, Kopecky KJ, et al. Revised recommendations of the International Working Group for Diagnosis, Standardization of Response Criteria, Treatment Outcomes and Reporting Standards for Therapeutic Trials in Acute Myeloid Leukemia. *J Clin Oncol*. 2003;21(24):4642-4649.
- Mantel N, Byar DP. Evaluation of response-time data involving transient states: an illustration using heart-transplant data. *J Am Stat Assoc*. 1974;69:81-86.
- Nucifora G, Larson RA, Rowley JD. Persistence of the 8;21 translocation in patients with acute myeloid leukemia type M2 in long term remission. *Blood*. 1993;82(3):712-715.
- Jurlander J, Caligiuri MA, Ruutu T, et al. Persistence of AML1-ETO fusion transcript in patients treated with allogeneic bone marrow transplantation for t(8;21) leukemia. *Blood*. 1996;88(6):2183-2191.
- Miyamoto T, Weissman IL, Akashi K. AML1/ETO-expressing non-leukemic stem cells in acute myelogenous leukaemia with 8;21 chromosome translocation. *Proc Natl Acad Sci U S A*. 2000;97(13):7521-7526.
- Tobal K, Johnson PRE, Saunders MJ, et al. Detection of CBF beta/MYH11 transcripts in patients with inversion and other abnormalities of chromosome 16 at presentation and remission. *Br J Haematol*. 1995;91(1):104-108.
- Elmaagacli AH, Beelen DW, Kroll M, et al. Detection of CBF beta/MYH11 fusion transcripts in patients with inv(16) acute myeloid leukaemia after allogeneic bone marrow or peripheral blood progenitor cell transplantation. *Bone Marrow Transplant*. 1998;21:159-166.
- Liu Yin JA. Minimal residual disease in acute myeloid leukaemia. *Baillieres Best Pract Res Clin Haematol*. 2002;15(1):119-135.
- Borthakur G, Faderl S, Verstovsek S. Molecular response in core binding factor acute myelogenous leukaemia with fludarabine, cytarabine, G-CSF and gemtuzumab ozogamicin. *Blood*. 2008;112:676.
- Breems DA, Van Putten WLT, Huijens PC, et al. Prognostic index for adult patients with acute myeloid leukemia in first relapse. *J Clin Oncol*. 2005;23(9):1969-1978.
- Marcucci G, Caligiuri MA, Dohner H, et al. Quantification of CBF beta/MYH11 fusion transcripts by real-time RT-PCR in patients with inv(16) acute myeloid leukaemia. *Leukemia*. 2001;15(7):1072-1080.
- Martinelli G, Rondoni M, Buonamici S, et al. Molecular monitoring to identify a threshold of CBFβ/MYH11 transcript below which continuous complete remission of acute myeloid leukemia inv 16 is likely. *Haematologica*. 2004;89(4):495-497.
- Boeckx N, De Roover J, van der Velden, et al. Quantification of CBFβ-MYH11 fusion gene levels in paired peripheral blood and bone marrow samples by real-time PCR. *Leukemia*. 2005;19(11):1988-2022.
- Schnittger S, Kern W, Tschulik C, et al. Minimal residual disease levels assessed by NPM1 mutation specific RQ-PCR provide important prognostic information in AML. *Blood*. 2009;14(11):2220-2231.
- Ommen HB, Schnittger S, Jovanovic JV, et al. Strikingly different molecular relapse kinetics in NPM1 c, PML-RARA, RUNX1-RUNX1T1 and CBFβ-MYH11 acute myeloid leukemia. *Blood*. 2010;115(2):198-205.
- Paschka P, Marcucci G, Ruppert AS, et al. Adverse prognostic significance of KIT mutations in adult acute myeloid leukemia with inv(16) and t(8;21): a Cancer and Leukemia Group B Study. *J Clin Oncol*. 2011;29(24):3904-3911.