

Multilineage dysplasia was detected in 32 cases (25%), a frequency similar to that reported by Falini, and was associated with a higher proportion of normal karyotype (93% vs 60%; $P < .001$), lower leukocyte count at diagnosis ($32 \times 10^9/L$ vs $69 \times 10^9/L$; $P = .01$), and lower bone marrow infiltration (51% vs 72% blast cells, $P < .001$). Interestingly, the frequency of *NPM1* and *FLT3* internal tandem duplication (*FLT3*-ITD) mutations did not differ between patients with and without MLD (59% vs 50%, and 31% vs 38%, respectively). *NPM1* mutations were found in 68 patients (52%). MLD was observed in 19 patients (28%) with mutated *NPM1* and in 13 (21%) with wild-type *NPM1*. Outcomes in patients with mutated *NPM1* were similar for those with and without MLD; response rate was 95% and 85%, 5-year relapse incidence was $35\% \pm 26\%$ and $47\% \pm 16\%$, and 5-year survival was $56\% \pm 23\%$ and $46\% \pm 14\%$, respectively. In contrast in patients with wild-type *NPM1*, those patients with MLD showed an inferior response rate to induction chemotherapy (53% vs 85%; $P = .02$). When the analysis was restricted to younger patients (≤ 60 years) those with MLD showed a lower 5-year survival (0% vs $40\% \pm 16\%$, $P = .012$; Figure 1). The unfavorable prognostic value of MLD on response rate ($P = .034$; relative risk, 4.8; 95% confidence interval, 1.1-20) and survival ($P = .036$; hazard ratio = 2.5; 95% confidence interval, 1.1-6) was confirmed in a multivariate analysis.

These results confirm that, although dysplastic features are a common trait in *NPM1*-mutated AML, they do not confer a worse prognosis. Falini et al found that gene expression profiling did not identify any distinctive MLD-associated gene signature in the mutated *NPM1* cohort.⁶ The correlation found in the present study between an unfavorable outcome and dysplastic features in wild-type *NPM1* IR-AML patients leads us to suggest that a search for novel genetic or epigenetic markers in this AML subgroup might reveal a specific biologic identity.

In conclusion, the prognostic relevance of MLD in IR-AML might be dependent on *NPM1* mutational status. Whereas MLD predicts an adverse outcome in patients with wild-type *NPM1*, it lacks prognostic value in *NPM1*-mutated AML. Nonetheless, this observation requires further confirmation in a larger series of patients.

Marina Díaz-Beyá

Hematology Department, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Hospital Clínic, Barcelona, Spain

María Rozman

Hematology Department, Hematopathology Unit, IDIBAPS, Hospital Clínic, Barcelona, Spain

Marta Pratcorona

Hematology Department, Hematopathology Unit, IDIBAPS, Hospital Clínic, Barcelona, Spain

Montserrat Torredadell

Hematology Department, Hematopathology Unit, IDIBAPS, Hospital Clínic, Barcelona, Spain

Mireia Camós

Hematology Department, Hematopathology Unit, IDIBAPS, Hospital Clínic, Barcelona, Spain

Josep Ll. Aguilár

Hematopathology Unit, IDIBAPS, Hospital Clínic, Barcelona, Spain

Jordi Esteve

Hematology Department, IDIBAPS, Hospital Clínic, Barcelona, Spain

Acknowledgments: This work has been partially supported by grants FIS03/0423, PI080158, and RD06/0020/0004 from "Instituto de Salud Carlos III" (ISCIII), Spanish Ministry of Health, Spain.

Contribution: M.D.-B. updated the database of patients included in the analysis, performed all statistical analysis, and wrote the manuscript; M.R. performed morphologic review of all cases and reviewed the article; M.P. updated the database, performed molecular analysis, and reviewed the article; M.T. and M.C. performed molecular analysis and reviewed the article; J.L.I.A. performed morphologic review of all cases and reviewed the article; and J.E. designed the study, supervised statistical analysis, and reviewed the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Dr Jordi Esteve, Hospital Clínic of Barcelona, Villarroel 170, Barcelona 08036, Spain; e-mail: jesteve@clinic.ub.es.

References

- Haferlach T, Schoch C, Löffler H, et al. Morphologic dysplasia in de novo acute myeloid leukemia (AML) is related to unfavorable cytogenetics but has no independent prognostic relevance under the conditions of intensive induction therapy: results of a multiparameter analysis from the German AML Cooperative Group studies. *J Clin Oncol*. 2003;21(2):256-265.
- Arber DA, Stein AS, Carter NH, Ikle D, Forman SJ, Slovak ML. Prognostic impact of acute myeloid leukemia classification. Importance of detection of recurring cytogenetic abnormalities and multilineage dysplasia on survival. *Am J Clin Pathol*. 2003;119(5):672-680.
- Wandt H, Schakel U, Kroschinsky F, et al. MLD according to the WHO classification in AML has no correlation with age and no independent prognostic relevance as analyzed in 1766 patients. *Blood*. 2008;111(4):1855-1861.
- Weinberg OK, Seetharam M, Ren L, et al. Clinical characterization of acute myeloid leukemia with myelodysplasia-related changes as defined by the 2008 WHO classification system. *Blood*. 2009;113(9):1906-1908.
- Falini B, Maciejewski K, Weiss T, et al. Multilineage dysplasia has no impact on biologic, clinicopathologic, and prognostic features of AML with mutated nucleophosmin (*NPM1*). *Blood*. 2010;115(18):3776-3785.
- Arber DA, Brunning RD, Orazi A, et al. Acute myeloid leukemia with myelodysplastic changes. In: Swerdlow SH, Campo E, Harris NL, et al, eds. *WHO Classification of Tumors of Hematopoietic and Lymphoid Tissues*. Lyon, France: International Agency for Research on Cancer (IARC); 2008:124-126.

To the editor:

Platelet secretion defect in patients with familial hemophagocytic lymphohistiocytosis type 5 (FHL-5)

Familial hemophagocytic lymphohistiocytosis (FHL) is a genetic disorder of lymphocyte cytotoxicity caused by mutations in the gene encoding perforin (FHL-2) or in genes encoding proteins

important for intracellular trafficking and exocytosis of perforin-containing lytic granules.¹ These include Munc13-4 (FHL-3), syntaxin 11 (FHL-4), and Munc18-2 (FHL-5). The molecular




Figure 1. Thrombin-induced granule secretion of FHL-5 platelets. Thrombin-mediated expression of CD62P (A) and CD63 (B) agonist on platelets was detected by flow cytometry. Diluted PRP ($5 \times 10^7/\text{mL}$) was stimulated with different concentrations of thrombin (0.05–1.0 U/mL; Dade Behring) in the presence of 1.25mM Gly-Pro-Arg-Pro/GPRP. Platelets were stained by monoclonal anti-CD62P antibody (CLBthromb/6-FITC, Immunotech) and anti-CD63 antibody (CLB-gran/12-FITC), respectively. Data are expressed as median of logarithmic fluorescence intensities.

machinery for the transport of lytic granules in part overlaps with that required for the transport of lysosome-related organelles in other cellular systems. This is well illustrated by immunodeficiency syndromes associated with albinism such as Griscelli syndrome type II (GSII), Chediak-Higashi syndrome (CHS), and Hermansky-Pudlak syndrome type 2 (HPS2). The genes affected in these diseases (*RAB27A*, *LYST*, and *AP3B1*) are also relevant for hair pigmentation, neurologic development, neutrophil development, and platelet function leading to manifestations such as partial albinism, mental retardation, neutropenia, and bleeding in addition to a high risk of hemophagocytic lymphohistiocytosis (HLH).^{2,3}

The recently described FHL-5 is considered to be mainly a disorder of cytotoxicity.^{4,5} A role for Munc18-2 has been demonstrated in mast cell degranulation,⁶ but a clinically relevant role for Munc18-2 deficiency in cell systems other than lymphocytes has not been documented. Munc18-2 is expressed in platelets and associates with a large complex containing synaptosome-associated protein of 23 kDa (SNAP-23) and cellubrevin/vesicle-associated membrane protein 3 (VAMP3).⁷ These data suggest that Munc18-2 might play a role in platelet granule exocytosis. In support of this concept, it was recently reported that 3 of 11 FHL-5 patients developed significant bleeding symptoms even outside of acute HLH episodes.⁸ However, all of these patients were also thrombocytopenic and platelet function tests were not performed.

Here we report on platelet function analysis of 4 patients with genetically confirmed FHL-5. The *MUNC18-2* mutations of these patients, their clinical course, and data showing impaired T cell and NK cell degranulation have been described previously^{4,9}: patients P1155 and P1945⁴ showed a typical course of FHL with manifestation of hemophagocytic syndrome

within the first year of life, while patients P1 and P4⁹ presented with a late onset of FHL. Of note, none of the patients had obvious bleeding symptoms.

Flow cytometric analyses of the patients' platelets revealed that platelet α (CD 62P)- and δ (CD 63)-granule secretion in response to thrombin stimulation was severely impaired in all patients (Figure 1). Surface expression of glycoprotein (GP) Ib/V/IX and GPIIb/IIIa, ristocetin-induced binding of Von Willebrand factor, and binding of soluble fibrinogen were normal in all patients. Platelet aggregation/agglutination after stimulation with ADP, collagen, and ristocetin was slightly impaired. Bleeding time was assessed in P1945 and was slightly prolonged (8.5 minutes). These data demonstrate a selective impairment of platelet granule secretion in patients with FHL-5 and thus document an important role for Munc18-2 in platelet degranulation. Platelet secretion defects have also been observed in patients with HPS2 or with CHS who can present with mucocutaneous bleedings, especially after surgery.^{2,10} Although bleeding symptoms in FHL-5 patients seem to be mild, our findings clearly demonstrate that Munc18-2 deficiency is more than a genetic disorder of cytotoxicity.

Kirstin Sandrock

Department of Pediatrics and Adolescent Medicine, University of Freiburg, Freiburg, Germany

Lea Nakamura

Department of Pediatrics and Adolescent Medicine, University of Freiburg, Freiburg, Germany

Thomas Vraetz

Centre of Chronic Immunodeficiency, University of Freiburg, Freiburg, Germany

Karin Beutel

Department of Pediatric Hematology and Oncology, University Medical Center, Hamburg-Eppendorf, Germany

Stephan Ehli

Centre of Chronic Immunodeficiency, University of Freiburg, Freiburg, Germany

Barbara Zieger

Department of Pediatrics and Adolescent Medicine, University of Freiburg, Freiburg, Germany

Acknowledgment: This work was supported by the German Federal Ministry of Education and Research (BMBF 01 EO 0803).

Contribution: L.N. performed research and analyzed data; K.S., S.E., and B.Z. designed research, analyzed data and wrote paper; and T.V. and K.B. took care of the patients. K.S. and L.N. contributed equally to this study.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Prof Dr Barbara Zieger, Department of Pediatrics and Adolescent Medicine, University of Freiburg, Mathildenstr 1, 79106 Freiburg, Germany; e-mail: barbara.zieger@uniklinik-freiburg.de.

References

- Schmid JP, Cote M, Menager MM, et al. Inherited defects in lymphocyte cytotoxic activity. *Immunol Rev*. 2010;235(1):10-23.
- Enders A, Zieger B, Schwarz K, et al. Lethal hemophagocytic lymphohistiocytosis in Hermansky-Pudlak syndrome type II. *Blood*. 2006;108(1):81-87.
- Stinchcombe J, Bossi G, Griffiths GM. Linking albinism and immunity: the secrets of secretory lysosomes. *Science*. 2004;305(5680):55-59.
- zur Stadt Rohr J, Seifert W, et al. Familial hemophagocytic lymphohistiocytosis type 5 (FHL-5) is caused by mutations in Munc18-2 and impaired binding to syntaxin 11. *Am J Hum Genet*. 2009;85(4):482-492.

- Cote M, Menager MM, Burgess A, et al. Munc18-2 deficiency causes familial hemophagocytic lymphohistiocytosis type 5 and impairs cytotoxic granule exocytosis in patient NK cells. *J Clin Invest*. 2009;119(12):3765-3773.
- Martin-Verdeaux S, Pombo I, Iannascoli B, et al. Evidence of a role for Munc18-2 and microtubules in mast cell granule exocytosis. *J Cell Sci*. 2003; 116(pt 2):325-334.
- Houng A, Polgar J, Reed GL. Munc18-syntaxin complexes and exocytosis in human platelets. *J Biol Chem*. 2003;278(22):19627-19633.
- Meeths M, Entesarian M, Al-Herz W, et al. Spectrum of clinical presentations in familial hemophagocytic lymphohistiocytosis (FHL) type 5 patients with mutations in STXBP2. *Blood*. 2010;116(15):2635-2643.
- Rohr J, Beutel K, Maul-Pavicic A, et al. Atypical familial hemophagocytic lymphohistiocytosis due to mutations in UNC13D and STXBP2 overlaps with primary immunodeficiency diseases. *Haematologica*. 2010;95(12): 2080-2087.
- Nurden P, Nurden AT. Congenital disorders associated with platelet dysfunctions. *Thromb Haemost*. 2008;99(2):253-263.

To the editor:

Platelet interior imaging technologies

Harry F. G. Heijnen and his colleagues recently reported studies of the platelet interior using electron tomography.¹ The methods used to freeze platelets and prepare them for cryo-electron tomography are of interest. However, most of the information presented is not new. A major finding was that the open canalicular system and dense tubular systems of channels were highly intertwined and formed close associations in specialized membrane regions. This observation is not new. Interaction of the 2 channel systems was first reported in 1972,² and the name “membrane complexes” assigned to them. Their similarity to the sarcoplasmic reticulum of embryonic muscle and ability of the dense tubular system to bind divalent cations suggested a role of membrane complexes in platelet muscle physiology.³

A review chapter, “Platelet Structure,” at page 70 (Figures 3-74, 3-75, and 3-76) in the second edition of Michelson’s textbook *Platelets*⁴ provides a more comprehensive picture of membrane complexes. Identification of α granule subtypes included organelles containing the multimeric von Willebrand factor assemblies in tubules eccentrically located in the α granule matrix.¹ Heijnen and colleagues cite a paper by Cramer et al⁵ for recognizing the tubular elements in α granules, demonstrating that they are von Willebrand factor multimers similar to Weibel-Palade bodies. However, an earlier study in 1968⁶ described them in thin platelet sections by transmission electron microscopy (TEM). The authors of the current paper¹ frequently encountered α granules with an elongated, “tubular shape,” and their Figures 4 and 5 provide images of the tubular subtypes. However, images in Figure 3-49 on

page 62 of Michelson’s text⁴ provide an image of α granules with extensions taken by TEM that is clearer than provided by tomography. The extensions appear as dense as the granules from which they originate. Therefore, we referred to them as “rod-like,” rather than “tubular.” The presence of cross-striations in the rods also was noted in another study.⁷

Electron tomography or its interpretation may make errors in organelle identification. The tomographic slices of a “tubular” α granule in Figure 5C-F and its reconstruction in 5G-H are interesting, but the organelle is not an α granule. It is a δ granule (dense body). Several examples are shown in Michelson⁴ on page 62 in Figures 3-50 and 3-51, on page 63 in Figure 3-52, and in examples included here (Figure 1). The images in the enclosed illustrations were taken by TEM on whole-mount preparations of normal platelets.

In summary, Heijnen et al have used a useful new technology to review the platelet interior. However, based on the concerns raised in this letter, it does not appear that electron tomography has replaced thin-section and whole-mount TEM.

James G. White
Regents Professor of Laboratory Medicine,
Pathology and Pediatrics,
University of Minnesota,
Minneapolis, MN

Conflict-of-interest disclosure: The author declares no competing financial interests. **Correspondence:** James G. White, MD, University of Minnesota




Figure 1. Whole-mounted platelets viewed by TEM. Two micrographs (A-B) of whole-mounted platelets viewed by TEM contain many spherical dense bodies (DBs) and DBs with 1 or 2 tails.